

AD-A141 708

DYNAMIC CHANGES OF PHASE IN A VAN DER WAALS FLUID(U)
WISCONSIN UNIV-MADISON MATHEMATICS RESEARCH CENTER
R HAGAN ET AL. MAR 84 MRC-TSR-2666 DAAG29-80-C-0041


1/1

UNCLASSIFIED

F/G 20/4

NL

END
DATE
FILMED
7 84
DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

2

MRC Technical Summary Report #2666

DYNAMIC CHANGES OF PHASE IN A
VAN DER WAALS FLUID

R. Hagan and J. Serrin

AD-A141 708

**Mathematics Research Center
University of Wisconsin—Madison
610 Walnut Street
Madison, Wisconsin 53705**

March 1984

(Received September 7, 1983)

DTIC FILE COPY

Approved for public release
Distribution unlimited

Sponsored by

U. S. Army Research Office
P. O. Box 12211
Research Triangle Park
North Carolina 27709

DTIC
ELECTE
JUN 01 1984
S D E

84 05 31 068

UNIVERSITY OF WISCONSIN-MADISON
MATHEMATICS RESEARCH CENTER

DYNAMIC CHANGES OF PHASE IN A VAN DER WAALS FLUID

R. Hagan and J. Serrin

Technical Summary Report #2666
March 1984

ABSTRACT

This paper gives sufficient conditions to guarantee the existence of a shock layer solution connecting two different equilibrium states in a van der Waals fluid. In particular, the equilibrium states can belong to two different phases of the fluid. The constitutive laws come from a modified Korteweg theory which is compatible with the Clausius Duhem inequality. The Clausius Duhem inequality in turn gives rise to a Liapunov function. The main mathematical tool is the LaSalle invariance principle.

AMS (MOS) Subject Classifications: 80A10, 76T05, 76N99, 35B99, 35M05,
35L65, 35L67

Key Words: phase transitions, shock layer, van der Waals fluid,
interstitial working, Liapunov function

Work Unit Number 2 (Physical Mathematics)

SIGNIFICANCE AND EXPLANATION

In the study of both static and dynamic phase transitions a typical model has the constitutive law for stress given by Korteweg's theory. In this theory, the classical form of Navier and Stokes has additional terms which model the effects of interfacial capillarity and depend on the spatial gradients of the density up to second order. It is these capillarity terms which allow for smooth static phase transitions. Unfortunately, Korteweg's theory is not compatible with the usual continuum theory of thermodynamics since the Clausius-Duhem inequality is, in general, not satisfied. Recently, a modified Korteweg theory has been developed by E. Dunn and J. Serrin, that does not suffer from this defect. We use this theory to generate a set of shock layer equations which are appropriate for shocks, including dynamic phase transitions. One of the immediate benefits obtained from this theory is that the Clausius-Duhem inequality gives rise to a Liapunov function for the shock layer equations. We then give sufficient conditions for the existence of a shock layer connecting two equilibrium points.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the authors of this report.

DYNAMIC CHANGES OF PHASE IN A VAN DER WAALS FLUID

R. Hagan and J. Serrin

INTRODUCTION

In a paper by Serrin [1], Korteweg's theory of capillarity [2,3] was used to find conditions for equilibrium between liquid and vapor phases of a van der Waals fluid. In a subsequent paper, Slemrod [4] extended Serrin's approach to study dynamic changes of phase in a van der Waals fluid, under the assumption of isothermal motion. This study was further extended by Hagan and Slemrod in [5]. The next logical step was to drop the assumption of isothermal motion. This was done for a van der Waals fluid in a paper [6] by Slemrod. In [6] he showed the existence of a shock layer that converts vapor to liquid and the existence of a shock layer that converts liquid to vapor, under assumptions that render the motion nearly isothermal. These assumptions are that the specific heat capacity at constant volume is large, the coefficients of heat conduction and viscosity are of the same small order μ , and the coefficients in the capillarity terms of the stress are of order μ^2 .

One of the problems that complicates the study of dynamic changes of phase is the incompatibility of the classical Korteweg stress with the Clausius Duhem inequality [7]. Recently, however, a modified Korteweg theory has been developed by Dunn and Serrin [8] that is compatible with the Clausius Duhem inequality. In this theory they posit the existence of a rate of supply of mechanical energy, the interstitial working, which takes into account the working of longer range interactions. With this additional term in the energy balance it is possible to derive a constitutive relation for stress that depends on spatial gradients of the density and still satisfies the Clausius Duhem inequality. These spatial gradient terms are used to model the effects of interfacial capillarity and at the same time allow the existence of static phase transitions [1], [9]. That is, if we were to use the classical form of Navier and Stokes for the stress then we would find that some dynamic phase transitions exist, but no static ones [10], [11].

In this paper we use a special form of the modified Korteweg theory contained in [8]. The Clausius Duhem inequality then gives a direct proof of the increase of entropy across a shock layer and it also provides a Liapunov function, in the sense of LaSalle [12], for the shock layer equations.

In section two of this paper we derive the shock layer equations and the increase of entropy theorem. In section three we examine some of the properties of the Hugoniot curves for a van der Waals fluid. In section four we state sufficient conditions to guarantee the existence of a compressive shock layer in a van der Waals fluid.

SECTION 2




Figure 1

We consider one-dimensional steady flow of a van der Waals fluid. The flow may be thought of as taking place in a cylinder of uniform cross-section parallel to the x -axis. We will always assume that the fluid velocity u is positive so that the fluid flows from left to right. The absolute temperature will be denoted by θ , and the specific volume by v .

We will seek a smooth solution $(u(x), v(x), \theta(x))$, $x \in \mathbb{R}$ of the equations of motion satisfying the following boundary conditions,

$$(2.1) \quad (u(x), v(x), \theta(x)) \rightarrow \begin{cases} (u_0, v_0, \theta_0) & \text{as } x \rightarrow -\infty \\ (u_1, v_1, \theta_1) & \text{as } x \rightarrow +\infty \end{cases},$$

$(u(x)', v(x)', \theta(x)') \rightarrow (0, 0, 0)$ as $x \rightarrow \pm\infty$, and $v''(x) \rightarrow 0$ as $x \rightarrow \pm\infty$. If such a smooth solution exists, it is called a shock layer. Of particular interest to us in this paper, is the question of the existence of a shock layer, when (v_0, θ_0) and (v_1, θ_1) belong to different phases of a fluid. If such a shock layer exists it will be called a dynamic phase transition.

The balance laws for mass, momentum, and energy for a one-dimensional steady flow, the absence of external body forces and radiant heating are (see [8])

$$(2.2) \quad (v^{-1}u)' = 0,$$

$$(2.3) \quad v^{-1}uu' = T'_{xx},$$

$$(2.4) \quad v^{-1}ue' = T'_{xx}u' + k' + q',$$

where $()' = d()/dx$. In addition to the balance laws we also have the Clausius Duhem inequality,

$$(2.5) \quad v^{-1}u\eta' \geq \left(\frac{q}{\theta}\right)'.$$

Here, T_{xx} is the x component of stress in the x direction, e is the specific internal energy, η is the specific entropy, k is the interstitial working, and q is the heat flux.

We will have need of the following notation, if f is a function of v and θ , then $f_1 = f(v_1, \theta_1)$, $f_0 = f(v_0, \theta_0)$, and $[f] = f_1 - f_0$.

Theorem 1.1. (The increase of entropy across a shock layer.)

Suppose $q = 0$ whenever $\theta' = 0$. Then $\eta_1 > \eta_0$ for any shock layer.

Proof.

Integration of (2.2) gives

$$(2.7) \quad u(x) = mv(x),$$

for some constant m . Furthermore $m > 0$ since $v > 0$, and $u > 0$ by assumption. We may now integrate (2.5) from $x = -\infty$ to $x = +\infty$, to obtain,

$$m(\eta_1 - \eta_0) \geq \frac{q}{\theta} \Big|_{-\infty}^{+\infty}.$$

But $\theta' = 0$ at $x = \pm\infty$, hence

$$(2.8) \quad \eta_1 > \eta_0.$$

In addition to the balance laws and the Clausius Duhem inequality we need to specify the constitutive structure of the fluid. We shall use the following special form of a modified Korteweg theory developed by J. E. Dunn and J. Serrin [8]. The two main features of this theory that makes it useful for work in phase transitions are the preservation of the Clausius Duhem inequality and the occurrence of higher spatial derivatives in the constitutive relation for stress.

The constitutive relations are (specialized to one-dimensional flow from equations (1.26) and (1.27) of [8])

$$(2.9) \quad k = \frac{\sigma u(v')^2}{v},$$

$$(2.10) \quad T_{xx} = -p + (\lambda + 2\mu)u' - \frac{\partial \sigma}{\partial v} \frac{(v')^2}{2} - \frac{\partial \sigma}{\partial \theta} v'\theta' - \sigma v'',$$

$$(2.11) \quad q = \kappa \theta',$$

$$(2.12) \quad \psi = \tilde{\psi} + \frac{\sigma}{2} (v')^2, \quad \text{the specific Helmholtz free energy,}$$

$$(2.13) \quad \eta = \tilde{\eta} - \frac{1}{2} \frac{\partial \sigma}{\partial \theta} (v')^2,$$

$$(2.14) \quad e = \tilde{e} + \frac{1}{2} \left(\sigma - \theta \frac{\partial \sigma}{\partial \theta} \right) (v')^2,$$

where $\sigma = cv^{-3}$ and c is the surface tension coefficient. Here $\tilde{\eta}$ and \tilde{e} are the equilibrium entropy and internal energy. They are functions of v and θ only, as is the pressure p . Furthermore, these quantities satisfy the classical Maxwell relationships of thermodynamics. In general σ , λ , μ , κ are positive continuous functions of v and θ , while σ , p , \tilde{e} , and $\tilde{\eta}$ are C^1 .

In this paper we shall restrict ourselves to the case where p and \tilde{e} satisfy the following hypotheses,

$$(H1) \quad p(v, \theta) = \frac{R\theta}{v-b} - \frac{a}{v^2}, \quad \text{for all } (v, \theta) \in \Omega,$$

$$(H2) \quad c_v = \frac{\partial \tilde{e}}{\partial \theta}(v, \theta) > 0, \quad \text{for all } (v, \theta) \in \Omega,$$

where Ω is defined as $\{(v, \theta) : b < v < \infty, 0 < \theta < \infty\}$. Here R, b , and a are positive constants and c_v is the specific heat at constant volume. (H1) is of course the van der Waals equation of state.

Let us substitute (2.7) into (2.3) and (2.4) to obtain,

$$(2.15) \quad mu' = T'_{xx} ,$$

$$(2.16) \quad me' = (T_{xx}u)' - T'_{xx}u + k' + q' .$$

We can now put (2.15) in (2.16) to obtain,

$$(2.17) \quad m(e + \frac{1}{2} u^2)' = (T_{xx}u)' + k' + q' .$$

Now we may integrate (2.15) and (2.17) from $-\infty$ to x and apply the boundary condition

(2.1). This gives

$$(2.18) \quad m(u - u_0) = T_{xx} - T_{xx_0} ,$$

$$(2.19) \quad m(e - \tilde{e}_0 + \frac{1}{2} (u^2 - u_0^2)) = T_{xx}u - T_{xx_0}u_0 + k + q .$$

We now insert (2.7), (2.9), (2.10), (2.11) and (2.14) into (2.18) and (2.19) to obtain,

$$(2.20) \quad \sigma v'' + \frac{\partial \sigma}{\partial \theta} v' \theta' + \frac{\partial \sigma}{\partial v} \frac{(v')^2}{2} - m(\lambda + 2\mu)v' + p - p_0 + m^2(v - v_0) = 0 ,$$

and

$$(2.21) \quad \kappa \theta' = m \left\{ -\frac{1}{2} \frac{\partial \sigma}{\partial \theta} (v')^2 + \tilde{e} - \tilde{e}_0 + p_0(v - v_0) - \frac{m^2}{2} (v - v_0)^2 \right\} .$$

It is convenient for us to define the following two functions,

$$(2.22) \quad L(v, \theta) = p(v, \theta) - p_0 + m^2(v - v_0) ,$$

and

$$(2.23) \quad M(v, \theta) = \tilde{e}(v, \theta) - \tilde{e}_0 + p_0(v - v_0) - \frac{1}{2} m^2(v - v_0)^2 .$$

We may now write (2.20) and (2.21) as a system of three first order ordinary differential equations, namely

$$(2.24a) \quad v' = w ,$$

$$(2.24b) \quad \sigma w' + \frac{\partial \sigma}{\partial \theta} w \theta' + \frac{\partial \sigma}{\partial v} \frac{w^2}{2} - m(\lambda + 2\mu)w + L(v, \theta) = 0 ,$$

$$(2.24c) \quad \kappa \theta' = -\frac{m}{2} \frac{\partial \sigma}{\partial \theta} w^2 + M(v, \theta), \quad (v, \theta) \in \Omega .$$

We shall refer to this system as the shock layer equations.

Lemma 1.2.

Given (u_0, v_0, θ_0) and (u_1, v_1, θ_1) with $(v_0, \theta_0), (v_1, \theta_1) \in \Omega$, a shock layer exists satisfying (2.1) if and only if there exist a solution to (2.24) satisfying,

$$(2.25) \quad (v(x), w(x), \theta(x)) \rightarrow \begin{cases} (v_0, 0, \theta_0) & \text{as } x \rightarrow -\infty \\ (v_1, 0, \theta_1) & \text{as } x \rightarrow \infty, \end{cases}$$

and (2.7) is satisfied.

Lemma 1.3. (The Rankine-Hugoniot jump conditions).

A necessary condition for a shock layer to exist satisfying (2.1) is that the Rankine-Hugoniot jump conditions are satisfied:

$$(2.26a) \quad [u] = m[v],$$

$$(2.26b) \quad [p] + m^2[v] = 0,$$

$$(2.26c) \quad [\tilde{e}] + \frac{p_1 + p_0}{2} [v] = 0.$$

Proof.

If a shock layer exists satisfying (2.1) then there exists a solution of (2.24) satisfying (2.25) and (2.7). Let $x \rightarrow \infty$; then we have from (2.24b) and (2.24c)

$$(2.27) \quad M_1 \approx L_1 = 0;$$

hence $[p] + m^2[v] = 0$ and $[\tilde{e}] + p_0[v] - \frac{1}{2} m^2[v]^2 = 0$. Therefore

$$[\tilde{e}] + \frac{1}{2} (p_1 + p_0)[v] = 0. \text{ Now let } x \rightarrow \pm\infty \text{ in (2.7); then } u_1 = mv_1 \text{ and } u_0 = mv_0.$$

Therefore $[u] = m[v]$.

We can solve the van der Waals equation of state for θ in terms of v and p algebraically. Hence we can define $\hat{e}(v, p) = \tilde{e}(v, \theta)$. We may then define

$$(2.28) \quad H(v, p; v_0, p_0) = \hat{e}(v, p) - \hat{e}(v_0, p_0) + \frac{p + p_0}{2} (v - v_0).$$

The curve in the $v - p$ plane consisting of all (v, p) satisfying $H(v, p; v_0, p_0) = 0$ is called the Hugoniot curve generated by (v_0, p_0) . Let us note that any state (v_1, p_1) lying in the intersection of the Hugoniot curve with the straight line given by

$$p = p_0 - m^2(v - v_0) \text{ will satisfy (2.26b, c) or equivalently (2.27). The corresponding}$$

jump $[u]$ is then given directly by (2.26a). Thus all conditions of (2.26) are satisfied.

It is convenient at this point to group together several thermodynamic identities which will be useful in the following sections. First, we have the standard Gibbs identity

$$(2.29) \quad \theta d\tilde{\eta} = d\tilde{e} + p dv$$

and the Maxwell relations

$$(2.30) \quad \frac{\partial \tilde{e}}{\partial v} = \theta \frac{\partial p}{\partial \theta} - p$$

$$(2.31) \quad \frac{\partial \tilde{\eta}}{\partial v} = \frac{\partial p}{\partial \theta}.$$

In addition, we shall need the formula

$$(2.32) \quad \left. \frac{\partial p}{\partial v} \right|_{\tilde{\eta}} = \frac{\partial p}{\partial v} - \frac{\theta}{c_v} \left(\frac{\partial p}{\partial \theta} \right)^2,$$

which follows from the chain rule

$$\frac{\partial p}{\partial v} = \left. \frac{\partial p}{\partial v} \right|_{\tilde{\eta}} + \left. \frac{\partial p}{\partial \tilde{\eta}} \right|_v \frac{\partial \tilde{\eta}}{\partial v}$$

together with (2.31) and the relation

$$\left. \frac{\partial p}{\partial \tilde{\eta}} \right|_v = \frac{\partial p}{\partial \theta} / \frac{\partial \tilde{\eta}}{\partial \theta} = \frac{\theta}{c_v} \frac{\partial p}{\partial \theta}.$$

From (2.30) one also gets $\partial \tilde{e} / \partial v = a/v^2$ for the van der Waals equation of state, so that in this case

$$(2.33) \quad \tilde{e} = -\frac{a}{v} + \int c_v(\theta) d\theta.$$

SECTION 3

The van der Waals equation of state possesses non-monotone isotherms for $\theta < \theta_c$, where θ_c is the critical temperature and monotone decreasing isotherms for $\theta > \theta_c$. (See Figure 2). We will define the unstable region in the $v - \theta$ plane as

$$(3.1) \quad \Omega_u = \{(v, \theta) \in \Omega : \frac{\partial p}{\partial v}(v, \theta) > 0\}.$$

Thus Ω_u is bounded by

$$(3.2) \quad \theta = \theta_u(v) \equiv \frac{2a}{R} \frac{(v-b)^2}{v^3}, \quad b < v < \infty.$$

We have $\frac{d\theta_u(v)}{dv} = \frac{-2a(v-b)(v-3b)}{Rv^4}$ and hence $\theta_u(v)$ has only one stationary point $v_c = 3b$ on (b, ∞) . Clearly, $\theta_u(v)$ takes on its maximum value $\theta_c = 8a/27Rb$, at $v = v_c$.




Figure 2. Isotherms for a van der Waals fluid.

We now define

$$(3.3) \quad \Omega_s = \{(v, \theta) \in \Omega : \theta_c < \theta < \infty\}$$

$$(3.4) \quad \Omega_l = \{(v, \theta) \in \Omega : \theta_u(v) < \theta < \theta_c, b < v < v_c\}$$

$$(3.5) \quad \Omega_u = \{(v, \theta) \in \Omega : \theta_u(v) < \theta < \theta_c, v_c < v < \infty\}$$

Here Ω_s is the super-heated vapor region, Ω_l is the liquid region, and Ω_u is the vapor region. (See Figure 3).




Figure 3

We shall on occasion refer to the fluid state in terms of v and p instead of v and θ . On such occasions it is useful to define the notation

$$\hat{\Omega}_\chi = \{(v, p(v, \theta)) : (v, \theta) \in \Omega_\chi\} \text{ where } \chi \text{ is } l, s, u, u, \text{ or empty.}$$

Note that the map $(v, \theta) \mapsto (v, p(v, \theta))$ is a homeomorphism of Ω onto $\hat{\Omega}$.

We now state and prove some useful lemmas concerning the Hugoniot curve.

Lemma 3.1.

Let $(v_0, p_0) \in \hat{\Omega}$ and let I be any compact interval such that $v_0 \in I \subseteq (b, \infty)$. Then the Hugoniot curve generated by (v_0, p_0) approaches the isotherm passing through (v_0, p_0) uniformly on I as $\inf_{0 < \theta < \infty} c_v(\theta)$ approaches infinity.

Proof.

$$H(v, p; v_0, p_0) = \hat{e}(v, p) - \hat{e}_0 + \frac{p + p_0}{2} (v - v_0) .$$

Hence

$$\frac{\partial H}{\partial p}(v, p; v_0, p_0) = \frac{\partial \hat{e}}{\partial p} + \frac{v - v_0}{2} .$$

Put

$$\frac{\partial \hat{e}}{\partial p} = \frac{\partial \tilde{e}}{\partial \theta} / \frac{\partial p}{\partial \theta} = \frac{c_v(v - b)}{R}$$

and so

$$\frac{\partial H}{\partial p} = \frac{c_v(v - b)}{R} + \frac{v - v_0}{2} \rightarrow +\infty \text{ as } \inf_{0 < \theta < \infty} c_v(\theta) \rightarrow \infty .$$

Therefore we can solve $H(v, p; v_0, p_0) = 0$ for p as a function of v , say $p = h(v)$, if

$\inf_{0 < \theta < \infty} c_v(\theta)$ sufficiently large. Assuming that this is so, we have

$$\left(\frac{\partial \hat{e}}{\partial p} + \frac{v - v_0}{2} \right) \frac{dh}{dv} + \frac{\partial \hat{e}}{\partial v} + \frac{p + p_0}{2} = 0 .$$

We also have

$$\frac{\partial \hat{e}}{\partial v} = \frac{\partial \tilde{e}}{\partial v} + \frac{\partial \tilde{e}}{\partial \theta} \frac{\partial \theta}{\partial v} \Big|_p = \frac{a}{v} - \frac{c_v(v - b)}{R} \frac{\partial p}{\partial v} \Big|_\theta .$$

Thus

$$(3.6) \quad \frac{dh}{dv} = \left(\frac{c_v(v - b)}{R} \frac{\partial p}{\partial v} \Big|_\theta - \frac{a}{v} - \frac{p + p_0}{2} \right) / \left(\frac{v - v_0}{2} + \frac{c_v(v - b)}{R} \right) ,$$

and hence $\frac{dh}{dv} + \frac{\partial p}{\partial v} \Big|_\theta$ uniformly on I as $\inf_{0 < \theta < \infty} c_v(\theta) \rightarrow \infty$. Thus

$$h(v) - h(v_0) + \int_{v_0}^v p(z, \theta_0) dz = p(v, \theta_0) - p_0$$

uniformly on I as $\inf_{0 < \theta < \infty} c_v(\theta) \rightarrow \infty$. Therefore $h(v) + p(v, \theta_0)$ uniformly on I as

$\inf_{0 < \theta < \infty} c_v(\theta) \rightarrow \infty$, since $h(v_0) = p(v_0, \theta_0)$.

Let us define

$$(3.7) \quad \gamma = \frac{R + c_v}{c_v} .$$

Lemma 3.2.

If c_v is constant, then $H(v, p; v_0, p_0) = 0$ can be solved algebraically for p as a function of v .

$$(3.8) \quad p \left\{ \frac{(\gamma + 1)}{2} v - \frac{(\gamma - 1)}{2} v_0 - b \right\} + \frac{a}{v^2} \{ (2 - \gamma)v - b \} =$$

$$p_0 \left\{ \frac{(\gamma + 1)}{2} v_0 - \frac{(\gamma - 1)}{2} v - b \right\} + \frac{a}{v_0^2} \{ (2 - \gamma)v_0 - b \}.$$

Let $p = h(v)$ be the solution of $H(v, p; v_0, p_0) = 0$ then

$$(3.9) \quad (v - v_s) \left\{ h(v) + \frac{\gamma - 1}{\gamma + 1} p_0 \right\} = \frac{2a}{\gamma + 1} \left\{ \frac{2\gamma(v_0 - b)}{a(\gamma + 1)} p_0 + \frac{\gamma - 2}{v} + \frac{b}{v^2} - \frac{\gamma - 2}{v_0} - \frac{b}{v_0^2} \right\},$$

where

$$(3.10) \quad v_s = \frac{(\gamma - 1)v_0 + 2b}{\gamma + 1}.$$

Proof.

We have $p = \frac{R\theta}{v - b} - \frac{a}{v^2}$, and hence

$$\theta = \frac{(v - b)}{R} \left(p + \frac{a}{v^2} \right)$$

$$\tilde{e} = c_v \theta - \frac{a}{v} = \frac{v - b}{\gamma - 1} \left(p + \frac{a}{v^2} \right) - \frac{a}{v}.$$

Now $H(v, p; v_0, p_0) = \tilde{e} - \tilde{e}_0 + \frac{p + p_0}{2} (v - v_0)$, and so the results follow.

It is clear from (3.9) that $h(v)$ has exactly one singularity at, $v = v_s$. As $c_v \uparrow \infty$, then $\gamma \uparrow 1$ and $v_s \uparrow b$; as $c_v \downarrow 0$, then $\gamma \uparrow \infty$ and $v_s \uparrow v_0$.

Lemma 3.3.

Suppose c_v is constant. Then

$$(3.11) \quad \lim_{v \rightarrow \infty} h(v) = - \frac{\gamma - 1}{\gamma + 1} p_0.$$

Proof.

The result follows at once from formula (3.9).

In the next lemma we shall examine the behavior of $h(v)$ as v approaches v_s . To this end we define

$$(3.12) \quad \sigma = \operatorname{sgn}\left(\frac{b^2}{a} p_0 + \frac{\gamma + 1}{\gamma((\gamma - 1)\gamma + 2)^2} \left\{ (\gamma - 1)(\gamma - 2) + \frac{4(\gamma - 1)}{\gamma} + \frac{2}{\gamma^2} \right\}\right)$$

where $\gamma = \frac{v_0}{b}$.

Lemma 3.4.

Suppose c_v is constant and $(v_0, p_0) \in \hat{\Omega}$. Then

$$(3.13) \quad \lim_{v \rightarrow v_s} h(v) = \begin{cases} -\infty, & \text{if } \sigma = 1 \\ +\infty, & \text{if } \sigma = -1 \end{cases},$$

$$(3.14) \quad \lim_{v \rightarrow v_s} h(v) = \begin{cases} +\infty, & \text{if } \sigma = 1 \\ -\infty, & \text{if } \sigma = -1 \end{cases},$$

$$(3.15) \quad \lim_{v \rightarrow v_s} h(v) = \frac{-a}{\gamma + 1} \left\{ \left(\frac{\gamma - 1}{a} \right) p_0 + \frac{(\gamma - 2)}{v_s^2} + \frac{2b}{v_s^3} \right\}, \text{ if } \sigma = 0.$$

Lemma 3.5.

If c_v is constant and $\sigma = 1$ or 0 and $\gamma > 2$ or $v_0 < \frac{(2\gamma - 1)b}{(\gamma - 1)(2 - \gamma)}$, then

$$h(v) < -\frac{(\gamma - 1)}{(\gamma + 1)} p_0 \text{ for } b < v < v_s.$$

Proof.

Let

$$(3.16) \quad f(v) = \frac{2\gamma(v_0 - b)}{a(\gamma + 1)} p_0 + \frac{\gamma - 2}{v} + \frac{b}{v^2} - \frac{\gamma - 2}{v_0} - \frac{b}{v_0^2},$$

so that the right hand side of equation (3.9) is given by $\frac{2a}{\gamma + 1} f(v)$. Lemma 3.4 now follows since $\sigma = \operatorname{sign} f(v_s)$. To obtain Lemma 3.5, we have by hypothesis that $\sigma = 1$ or 0 and hence $f(v_s) > 0$. We may write (3.16) as,

$$(3.17) \quad f(v) = f(v_g) + \frac{\gamma - 2}{v} + \frac{b}{v^2} - \frac{\gamma - 2}{v_g} - \frac{b}{v_g^2}.$$

If $\gamma > 2$, then $f(v) > 0$ for $b < v < v_g$. Now suppose that $1 < \gamma < 2$. The first and second derivatives of $f(v)$ are,

$$f'(v) = \frac{2 - \gamma}{v^2} - \frac{2b}{v^3} \quad \text{and} \quad f''(v) = -2\left(\frac{2 - \gamma}{v^3} - \frac{3b}{v^4}\right).$$

Now $f'(v)$ has only one zero at $\bar{v} = \frac{2b}{2 - \gamma}$ and $f''(\bar{v}) = \frac{2b}{\bar{v}^4} > 0$. So that $f(v)$ takes on its minimum value at \bar{v} . If $v_g < \bar{v}$ then $f(v) > f(v_g) > 0$ for $b > v < v_g$ since $\sigma = \text{sgn } f(v_g) = 1$ or 0 and $f(v)$ strictly decreasing on $b < v < \bar{v}$. But $v_g < \bar{v}$ if and only if $v_0 < \frac{(2\gamma - 1)b}{(\gamma - 1)(2 - \gamma)}$ by definition. Thus under the hypotheses of the lemma $f(v) > 0$ for $b < v < v_g$. Now (3.9) can be written as

$$(3.18) \quad (v - v_g) \left\{ h(v) + \frac{\gamma - 1}{\gamma + 1} p_0 \right\} = \frac{2a}{\gamma + 1} f(v)$$

and thus

$$(3.19) \quad h(v) < -\frac{\gamma - 1}{\gamma + 1} p_0 \quad \text{for } b < v < v_g.$$

Lemma 3.6.

If the hypotheses of Lemma 3.5 are satisfied then the Rankine-Hugoniot jump conditions cannot be satisfied if $p_0 > 0$ and $v_1 < v_g$.

Proof.

In order for the Rankine-Hugoniot jump conditions to be satisfied, we must have

$$-m^2 = \frac{h(v_1) - p_0}{v_1 - v_0},$$

and so $h(v_1) - p_0 > 0$, since $v_1 < v_0$. But by Lemma 3.5 we have $h(v_1) - p_0 < \frac{-2\gamma p_0}{\gamma + 1} < 0$ and hence the conclusions follows.

SECTION 4.

Theorem 4.1.

Assume that $(v_1, p_1), (v_0, p_0) \in \hat{\Omega} \setminus \text{closure } (\hat{\Omega}_u)$ and that $v_1 < v_0$.
 $H(v_1, p_1; v_0, p_0) = 0$, and

$$\frac{p_1 - p_0}{v_1 - v_0} = -m^2 < 0.$$

Furthermore, suppose that the chord connecting (v_1, p_1) to (v_0, p_0) lies above the graph of $H(v, p; v_0, p_0) = 0$ on the interval $v_1 < v < v_0$ and is not tangent to the graph at either end point. Assume finally that the straight line extension of this chord does not intersect the graph of $H(v, p; v_0, p_0) = 0$ when $v > v_0$ and $p > p(v, 0)$. Then there exists a unique compression shock-layer connecting (v_0, p_0) to (v_1, p_1) . Furthermore, $\theta_1 > \theta_0$, $\tilde{n}_1 > \tilde{n}_0$, the flow is supersonic at the state (v_0, p_0) , and subsonic at the state (v_1, p_1) .

The proof of Theorem 4.1 will be carried out with the aid of the following lemmas.




Figure 4

Lemma 4.2.

Each of the equations $M(v, \theta) = 0$ and $L(v, \theta) = 0$ uniquely define θ as a function of v , say $\theta = \theta_M(v)$ and $\theta_L(v)$. Furthermore under the hypotheses of Theorem 4.1 the curve $L = 0$ intersects the v -axis in exactly two points $v = b$ and $v = \bar{v}$, where $\bar{v} > v_0$. Moreover the curve $L = 0$ lies above (below) the curve $M = 0$ in Ω when $v_1 < v < v_0$ ($v_0 < v < \bar{v}$). That is

$$(4.1) \quad \theta_L(v) > \theta_M(v) \quad \text{for } v_1 < v < v_0$$

$$(4.2) \quad \theta_L(v) < \theta_M(v) \quad \text{for } v_0 < v < \bar{v}.$$

Proof.

We have

$$(4.3) \quad \frac{\partial M}{\partial \theta} = c_v > 0$$

and

$$(4.4) \quad \frac{\partial L}{\partial \theta} = \frac{\partial p}{\partial \theta} > 0$$

so that $M = 0$ and $L = 0$ can be solved for θ as a function of v . In particular

$$(4.5) \quad \theta_L(v) = \frac{v - b}{R} \left(\frac{a}{v^2} + p_0 - m^2(v - v_0) \right).$$

It is clear from (4.5) that $\theta_L(b) = 0$. In the $v - p$ plane the line $L = 0$ intersects the lower boundary of the region $\hat{\Omega}$, given by $p = p(v, 0)$, $b < v < \infty$, only once since $L = 0$ has negative slope and $p = p(v, 0)$ has positive slope. Therefore in the $v - \theta$ plane the curve $L = 0$ intersects the v -axis (i.e. $\theta = 0$) only once in the interval (b, ∞) .

The curves $L = 0$ and $M = 0$ intersect in Ω if and only if the line $L = 0$ and the curve $M = 0$ intersect in $\hat{\Omega}$, as follows immediately from the relation

$$(4.6) \quad H = M + \frac{1}{2} (v - v_0) L.$$

Thus it is sufficient to prove that the curve $L = 0$ lies above the curve $M = 0$ for at least one value of v in the interval (v_1, v_0) . We have

$$(4.7) \quad \frac{\partial H}{\partial p} \Big|_v = \frac{c_v(v - b)}{R} + \frac{(v - v_0)}{2} > 0$$

in some neighborhood of (v_0, p_0) . Hence, by our assumption, in this neighborhood $H > 0$ on $L = 0$ if $v < v_0$. Thus by (4.6), in this neighborhood we also have $M > 0$ on $L = 0$ when $v < v_0$.

From Lemma 4.2 and equation (4.5) we see that the graph of $\theta_L(v)$ lies in Ω only when the domain of $\theta_L(v)$ is (b, \bar{v}) . In the next lemma we shall show that the graph of $\theta_M(v)$ lies in Ω when $b < v < \infty$.

Lemma 4.3.

Under the hypotheses of Theorem 4.1 there is a constant $\bar{\theta} > 0$ such that $\theta_M(v) > \bar{\theta}$ for all $v > b$, with equality holding only at $v = \bar{v}$.

Proof.

The curve $M = 0$ lies above the curve $L = 0$ for $v_0 < v < \bar{v}$ by Lemma 4.2, and the curve $L = 0$ lies above the v -axis for $v_0 < v < \bar{v}$. It is sufficient to show that the curve $\theta = \theta_M(v)$ takes on its minimum value in the interval (b, ∞) at $v = \bar{v}$. Now

$$(4.8) \quad \frac{\partial M}{\partial v} = \theta \frac{\partial p}{\partial \theta} - L$$

and by (4.3), since $c_v = c_v(\theta)$,

$$(4.9) \quad \frac{\partial^2 M}{\partial \theta \partial v} = 0$$

so that

$$(4.10) \quad \frac{\partial M}{\partial v}(v, \theta) = \frac{\partial M}{\partial v}(v, 0) = -L(v, 0).$$

From (4.5) and Lemma 4.2 we see that $L(v, 0) < 0$ when $b < v < \bar{v}$ and $L(v, 0) > 0$ when $\bar{v} < v < \infty$. But

$$(4.11) \quad \frac{d\theta_M(v)}{dv} = - \frac{\partial M}{\partial v}(v, 0) / \frac{\partial M}{\partial \theta}(v, \theta) = \frac{L(v, 0)}{c_v(\theta)}$$

and so $\theta_M(v)$ takes on its minimum at $v = \bar{v}$ and $\theta_M(v) > \theta_M(\bar{v})$ for all $v > b$, with equality holding only at $v = \bar{v}$.

Lemma 4.4.

Under the hypotheses of Theorem 4.1 the critical point $(v_1, 0, \theta_1)$ is a saddle point with a one-dimensional stable manifold and a two dimensional unstable manifold.

Furthermore the fluid velocity is subsonic at the back state (v_1, θ_1) and is supersonic at the front state (v_0, θ_0) .

Proof.

The acoustic speed c is given by

$$(4.12) \quad c^2 = \frac{\partial p}{\partial \rho} \Big|_{\tilde{\eta}} = -v^2 \frac{\partial p}{\partial v} \Big|_{\tilde{\eta}}.$$

Thus

$$(4.13) \quad u^2 - c^2 = v^2 \left(\frac{\partial p}{\partial v} \Big|_{\tilde{\eta}} + m^2 \right)$$

since $u = mv$. We need to show that

$$(4.14) \quad \frac{\partial p}{\partial v} \Big|_{\tilde{\eta}} + m^2 > 0 \quad \text{at } (v_0, \theta_0)$$

and

$$(4.15) \quad \frac{\partial p}{\partial v} \Big|_{\tilde{\eta}} + m^2 < 0 \quad \text{at } (v_1, \theta_1).$$

Now by hypotheses the chord connecting (v_1, θ_1) to (v_0, θ_0) has slope $-m^2$ and lies above the Hugoniot curve ($H = 0$ curve) in $\hat{\Omega}$. Thus the slope of the curve $H = 0$ is greater than the slope of the chord at (v_0, θ_0) .

From (2.28) and the Gibbs relation we have

$$(4.16) \quad dH = \left(\theta \frac{\partial \tilde{\eta}}{\partial p} \Big|_v + \frac{v - v_0}{2} \right) dp + \left(\theta \frac{\partial \tilde{\eta}}{\partial v} \Big|_p - \frac{p - p_0}{2} \right) dv.$$

Thus

$$(4.17) \quad \frac{dp}{dv} \Big|_H = - \frac{\partial \tilde{\eta}}{\partial v} \Big|_p / \frac{\partial \tilde{\eta}}{\partial p} \Big|_v = \frac{\partial p}{\partial v} \Big|_{\tilde{\eta}} \quad \text{at } (v_0, p_0)$$

and so

$$\frac{\partial p}{\partial v} \Big|_{\tilde{\eta}} + m^2 > 0 \quad \text{at } (v_0, p_0).$$

This proves (4.14). We next show (4.15).

By Lemma 4.2 the curve $L = 0$ lies below the curve $M = 0$ for $v_0 < v < \bar{v}$ and so

$$(4.18) \quad \frac{d\theta}{dv} \Big|_L (v_1) > \frac{d\theta}{dv} \Big|_M (v_1).$$

The strictness of the inequality can be shown to follow from the non-tangency hypothesis and equation (4.4). Now

$$(4.19) \quad \frac{d\theta}{dv} \Big|_L = - \frac{\partial L}{\partial v} / \frac{\partial L}{\partial \theta}$$

and

$$(4.20) \quad \frac{d\theta}{dv} = - \frac{\partial M}{\partial v} / \frac{\partial M}{\partial \theta}$$

thus by (4.18)

$$(4.21) \quad D \equiv \frac{\partial L}{\partial v} \frac{\partial M}{\partial \theta} - \frac{\partial M}{\partial v} \frac{\partial L}{\partial \theta} < 0 \quad \text{at } (v_1, \theta_1)$$

since $\frac{\partial L}{\partial \theta}$ and $\frac{\partial M}{\partial \theta}$ are positive. Substituting (4.3), (4.4), (4.8), and

$$(4.22) \quad \frac{\partial L}{\partial v} = \frac{\partial p}{\partial v} + m^2$$

into (4.21) gives

$$(4.23) \quad D = c_v \left(\frac{\partial p}{\partial v} + m^2 \right) - \left(\theta \frac{\partial p}{\partial \theta} - L \right) \frac{\partial p}{\partial \theta} \quad \text{at } (v_1, \theta_1).$$

But $L(v_1, \theta_1) = 0$ and (see Section 2)

$$(4.24) \quad \frac{\partial p}{\partial v} - \frac{\theta}{c_v} \left(\frac{\partial p}{\partial \theta} \right)^2 = \frac{\partial p}{\partial v} \Big|_{\tilde{\eta}}$$

so that

$$(4.25) \quad D = c_v \left(\frac{\partial p}{\partial v} \Big|_{\tilde{\eta}} + m^2 \right).$$

Since $D < 0$, condition (4.15) follows at once.

To show that $(v_1, 0, \theta_1)$ is a saddle point we linearize (2.24) about $(v_1, 0, \theta_1)$ to obtain

$$(4.26) \quad \begin{pmatrix} v' \\ w' \\ \theta' \end{pmatrix} = A \begin{pmatrix} v - v_1 \\ w \\ \theta - \theta_1 \end{pmatrix}$$

where

$$(4.27) \quad A = \begin{pmatrix} 0 & 1 & 0 \\ -\frac{1}{\sigma} \frac{\partial L}{\partial v} & m \frac{\lambda + 2\mu}{\sigma} & -\frac{1}{\sigma} \frac{\partial L}{\partial \theta} \\ \frac{m}{\kappa} \frac{\partial M}{\partial v} & 0 & \frac{m}{\kappa} \frac{\partial M}{\partial \theta} \end{pmatrix},$$

the entries being evaluated at (v_1, θ_1) .

Now at (v_1, θ_1)

$$(4.28) \quad \det A = \frac{m}{\kappa \sigma} \left(\frac{\partial L}{\partial v} \frac{\partial M}{\partial \theta} - \frac{\partial L}{\partial \theta} \frac{\partial M}{\partial v} \right) = \frac{m}{\kappa \sigma} D < 0$$

by (4.21). Also

$$(4.29) \quad \text{trace } A = m\left(\frac{\lambda + 2\mu}{\sigma} + \frac{c}{\kappa} \frac{v}{\kappa}\right) > 0.$$

Let ρ_1, ρ_2 and ρ_3 be the eigenvalues of A . Suppose first that these are all real and that $\rho_1 < \rho_2 < \rho_3$. Then $\rho_1 \rho_2 \rho_3 = \det A < 0$ by (4.28) and so either $\rho_1 < 0, \rho_2 < 0$, and $\rho_3 < 0$ or $\rho_1 < 0, \rho_2 > 0$, and $\rho_3 > 0$. But $\rho_1 + \rho_2 + \rho_3 = \text{trace } A > 0$ by (4.29), and hence $\rho_1 < 0, \rho_2 > 0$ and $\rho_3 > 0$. Next suppose that ρ_1, ρ_2 and ρ_3 are not all real. Let $\rho_1 = \alpha - i\beta$ and $\rho_2 = \alpha + i\beta$, where α, β and ρ_3 are all real. Then $(\alpha^2 + \beta^2)\rho_3 = \det A < 0$; hence $\rho_3 < 0$ and $2\alpha + \rho_3 = \text{trace } A > 0$, so $\alpha > 0$. Therefore the critical point $(v_1, 0, \theta_1)$ is a saddle point with a one-dimensional stable manifold and a two-dimensional unstable manifold.

Define

$$(4.30) \quad \Phi(v, w, \theta) = m(\tilde{\eta} - \tilde{\eta}_1 + \frac{\sigma w^2}{2\theta} - \frac{M}{\theta}).$$

Lemma 4.5.

Let $(v(x), w(x), \theta(x))$ be any solution of (2.24). Then

$$(4.31) \quad \frac{d\Phi}{dx}(v(x), w(x), \theta(x)) \geq 0.$$

That is Φ is nondecreasing along the trajectories of (2.24).

Proof.

We can use (2.13), (2.23), and (2.24c) to write (4.30) as

$$(4.32) \quad \Phi = m(\eta - \eta_1) - \frac{\kappa \theta'}{\theta}.$$

But

$$(4.33) \quad \Phi' = m\eta' - \left(\frac{\kappa \theta'}{\theta}\right)' > 0$$

by the Clausius-Duhem inequality, completing the proof.

Now $\Phi(v_1, 0, \theta_1) = 0$ and

$$(4.33) \quad \frac{\partial \Phi}{\partial \theta}(v, 0, \theta) = \frac{m}{\theta^2} M(v, \theta)$$

so that $\frac{\partial \Phi}{\partial \theta}(v_1, 0, \theta) > 0$ (< 0) for $\theta > \theta_1$ ($\theta < \theta_1$). Hence

$$(4.34) \quad \Phi(v_1, 0, \theta) > 0 \text{ for } \theta \neq \theta_1.$$

The function $\theta_L(v)$ has a maximum on the interval (b, \bar{v}) , say $\bar{\theta}$, since by (4.5) we see that $\theta_L(b) = \theta_L(\bar{v}) = 0$ and $\theta_L(v) > 0$ for $b < v < \bar{v}$. Now

$$(4.35) \quad \frac{\partial \Phi}{\partial v}(v, 0, \theta) = \frac{m}{\theta} L(v, \theta)$$

and so

$$(4.36) \quad \frac{\partial \Phi}{\partial v}(v, 0, \bar{\theta}) > 0$$

since $L > 0$ to the right of the line $v = \bar{v}$. Thus

$$(4.37) \quad \Phi(v, 0, \bar{\theta}) > 0 \quad \text{for } v > v_0$$

since $\Phi(v_1, 0, \bar{\theta}) > 0$. Equality occurs if $\theta_1 = \bar{\theta}$.

We next show that the curve $M = 0$ intersects the line $\theta = \bar{\theta}$ for some $\bar{v} > \bar{v}$. Now from (4.11) we have

$$(4.38) \quad \frac{d\theta}{dv} \frac{M}{v}(v) = \frac{L(v, 0)}{c} = \frac{1}{c} (m^2(v - v_0) - p_0 - \frac{a}{v^2})$$

and so by the definition of \bar{v} ,

$$(4.39) \quad \frac{d\theta}{dv} \frac{M}{v}(\bar{v}) = 0.$$

Therefore, $\theta_M(v) \uparrow \infty$ as $v \uparrow \infty$ since by (4.38) $\frac{d\theta}{dv} \frac{M}{v}(v) \uparrow \infty$ as $v \uparrow \infty$. Thus the curve $M = 0$ intersects the line $\theta = \bar{\theta}$ for some $\bar{v} > \bar{v}$. From (4.38) and (4.33) it is clear that $\Phi(\bar{v}, 0, \theta) > 0$ for $0 < \theta < \bar{\theta}$.

Put

$$(4.40) \quad \bar{w} = \sqrt{\max_{\substack{v_1 < v < \bar{v} \\ \bar{\theta} < \theta < \bar{\theta}}} -2\theta\Phi(v, 0, \theta)}$$

We define a box B in phase space by

$$(4.41) \quad B = \{(v, w, \theta); v_1 < v < \bar{v}, -\bar{w} < w < \bar{w}, \bar{\theta} < \theta < \bar{\theta}\}.$$

Note that with the exception of the bottom of the box and the point $(v_1, 0, \theta_1)$ we have

$\Phi > 0$ on ∂B .

Lemma 4.6.

Under the hypotheses of Theorem 4.1 every trajectory which intersects the bottom of the box B leaves the box.

Proof.

We have from (2.24c)

$$w\theta' = m\left(-\frac{1}{2} \frac{\partial \theta}{\partial v} w^2 + M(v, \theta)\right).$$

From Lemma 4.3 the curve $M = 0$ does not drop below the line $\theta = \bar{\theta}$ in Ω . But

$$\frac{\partial M}{\partial \theta} = c_v > 0 \text{ and so}$$

$$(4.42) \quad M(v, \theta) < 0 \text{ for } b < v < \infty \text{ and } 0 < \theta < \bar{\theta}$$

with equality holding only if $(v, \theta) = (\bar{v}, \bar{\theta})$. Now

$$(4.43) \quad \frac{\partial \sigma}{\partial \theta} = \sigma + \frac{\theta \partial \sigma}{\partial \theta}$$

and by hypothesis $\sigma > 0$ and

$$(4.44) \quad \liminf_{\theta \rightarrow 0} \frac{\theta \partial \sigma}{\partial \theta} (v, \theta) > 0 \text{ for } b < v < \infty$$

so that

$$(4.44) \quad \frac{\partial \sigma}{\partial \theta} > 0 \text{ for } b < v < \infty$$

and θ sufficiently small and positive. We can choose $\bar{\theta}$ smaller if necessary so that

(4.44) holds when $0 < \theta < \bar{\theta}$. Thus $\theta' < 0$ on the bottom of the box B with the possible exception of $(\bar{v}, 0, \bar{\theta})$. Therefore every trajectory which intersects the bottom of the box at some point leaves the box at that point.

Lemma 4.7.

Under the hypotheses of Theorem 4.1 one of the trajectories of the stable manifold of $(v_1, 0, \theta_1)$ enters the box B while the other never does. Thus there can be at most one trajectory of (2.24) connecting $(v_0, 0, \theta_0)$ to $(v_1, 0, \theta_1)$.

Proof.

We need to show that the line tangent to the stable manifold is transverse to the plane $v = v_1$ in phase space. This line is parallel to the eigenvector associated with the negative eigenvalue ρ of (4.27). Let $(\xi_1, \xi_2, \xi_3)^T$ be this eigenvector. Then we have

$$(4.44) \quad A \begin{pmatrix} \xi_1 \\ \xi_2 \\ \xi_3 \end{pmatrix} = \rho \begin{pmatrix} \xi_1 \\ \xi_2 \\ \xi_3 \end{pmatrix}$$

We assert that $\xi_1 \neq 0$, which is the required transversality condition. Suppose for contradiction that $\xi_1 = 0$. Then from (4.44) and (4.27) we have

$$(4.45) \quad \begin{aligned} \xi_2 &= \rho \xi_1 = 0 \\ (\kappa \rho - mc_v) \xi_3 &= m \frac{\partial M}{\partial v} \xi_1 = 0. \end{aligned}$$

But $\kappa \rho - mc_v < 0$. Hence $\xi_1 = \xi_2 = \xi_3 = 0$, which is impossible.

Now suppose one of the trajectories forming the stable manifold of $(v_1, 0, \theta_1)$ crosses the plane $v = v_1$ at some point. Then $\phi > 0$ at this point since

$$(4.46) \quad \phi(v, w, \theta) = \phi(v, 0, \theta) + \frac{m \omega w^2}{2\theta}.$$

and $\phi(v_1, 0, \theta) > 0$ for $\theta \neq \theta_1$ by (4.34). By Lemma 4.5, ϕ is nondecreasing along trajectories of (2.24), and so $\phi(v_1, 0, \theta_1) > 0$. But $\phi(v_1, 0, \theta_1) = 0$ and hence any trajectory forming the stable manifold cannot cross the plane $v = v_1$.

If we replace x by $-x$ in the system of ordinary differential equations (2.24) then the direction of each trajectory in (2.24) will be reversed. We shall refer to (2.24) with x replaced by $-x$ as the reversed system. Note that ϕ decreases along any trajectory of the reversed system and hence ϕ is a Liapunov function in the sense of LaSalle for the reversed system.

Let us denote by \emptyset the trajectory of the reversed system that leaves $(v_1, 0, \theta_1)$ and enters B .

Lemma 4.8.

Under the hypotheses of Theorem 4.1 the trajectory \emptyset is bounded.

Proof.

We shall show that \emptyset is contained in B . For the reversed system, ϕ is nonincreasing along \emptyset and hence $\phi \leq 0$ on \emptyset since $\phi(v_1, 0, \theta_1) = 0$. But $\phi > 0$ on ∂B with the exception of the point $(v_1, 0, \theta_1)$ and the bottom face of B . Thus \emptyset cannot cross ∂B except possibly at the bottom face. But by Lemma 4.6 and the fact that the system is reversed, every trajectory which intersects the bottom face enters B . Thus \emptyset cannot cross ∂B and so it is bounded.

To complete the proof of Theorem 4.1 we need to show that \bar{O} enters $(v_0, 0, \theta_0)$, that is $\omega(\bar{O}) = \{(v_0, 0, \theta_0)\}$ for the reversed system. Since Φ is a Liapunov function in the sense of LaSalle for the reversed system and \bar{O} is bounded, $\omega(\bar{O})$ is contained in the largest invariant subset of $S \equiv \{(v, w, \theta) \in \text{closure of } B : \Phi' = 0\}$, by LaSalle's invariance principle [12].

From (4.33) we have

$$(4.47) \quad \Phi' = m\eta' - \frac{(\kappa\theta')'}{\theta} + \frac{\kappa\theta'^2}{3^2}$$

Now upon substituting (2.24c) and (2.13) into (4.47) we have

$$(4.48) \quad \Phi' = \frac{m^2}{\theta} (\lambda + 2\mu)w^2 + \frac{\kappa\theta'^2}{\theta^2}.$$

Therefore $\Phi' = 0$ implies $w = \theta' = 0$ and by (2.24) also $\sigma w' + L = 0$ and $M = 0$ when $\Phi' = 0$. Thus the largest invariant subset of S is $\{(v_1, 0, \theta_1), (v_0, 0, \theta_0)\}$. But $\theta' \neq 0$ on \bar{O} so that $\Phi' < 0$ somewhere on \bar{O} and hence $\Phi < 0$ on \bar{O} . Therefore

$\omega(\bar{O}) = \{(v_0, 0, \theta_0)\}$ and so the trajectory \bar{O} connects $(v_1, 0, \theta_1)$ to $(v_0, 0, \theta_0)$.

It remains to show that $\tilde{\eta}_1 > \tilde{\eta}_0$ and $\theta_1 > \theta_0$. The first follows from (2.8), since (4.30) and (4.48) show that the equality cannot hold. The second follows from (4.11) since $L(v, 0) < 0$ for $v_1 < v < v_0$.

Example 1.

Suppose that c_v is a large constant, so that by Lemma 3.1 the Hugoniot curve generated by (v_0, p_0) is near the isotherm $p(v, \theta_0)$. Let us suppose that (v_0, p_0) is in the vapor region and (v_1, p_1) is in the liquid region. Furthermore, suppose that the straight line through (v_0, p_0) and (v_1, p_1) intersects the Hugoniot curve at only these two points (see Figure 5).

In this example the hypotheses of Theorem 4.1 are satisfied and hence there is a shock connecting (v_0, p_0) to (v_1, p_1) . The shock layers converts vapor in the equilibrium state (v_0, p_0) into liquid in the equilibrium state (v_1, p_1) . We shall call such a layer a liquefaction layer [13].




Figure 5

Example 2.

We keep the same set up as in Example 1, but with (v_1, p_1) also in the vapor region and $v_1 < v_0$ (see Figure 6). Then there is a gas-gas compressive shock layer converting gas in the more rarefied equilibrium state (v_0, p_0) to gas in the hotter denser equilibrium state (v_1, p_1) .




Figure 6

Example 3.

Let us suppose that c_v is constant and $0 < c_v < R$, so that $\gamma > 2$. Let us choose (v_0, p_0) in either the vapor or super-heated vapor region, with $v_0 > 7b$ and $p_0 > 0$. Then $\sigma = 1$ by (3.12), and thus the only reachable states on the Hugoniot curve are to the right of

$$v_s = \frac{(\gamma - 1)v_0 + 2b}{\gamma + 1}$$

by Lemma 3.6. Since $v_0 > 7b$ and $\gamma > 2$ we have

$$v_s > \frac{(\gamma - 1)7b + 2b}{\gamma + 1} = 7b - \frac{12b}{\gamma + 1} > 3b.$$

But the liquid region lies to the left of $v = 3b$, and hence no liquification shock is possible (see Figure 7).




Figure 7

REFERENCES

1. J. Serrin, Phase transitions and interfacial layers for van der Waals fluids. Recent Methods in Nonlinear Analysis and Applications, A. Camforo, S. Rionero, C. Sbordone, C. Trombetti, editors. Naples (1980).
2. D. J. Korteweg, Sur la forme que prennent les équations des mouvement des fluides si l'on tient compte des forces capillaires par des variation de densité, Archives Neerlandaises des Science Exactes et Naturelles, II 6, 1-24 (1901).
3. C. A. Truesdell, W. Noll, The non-linear field theories of mechanics, Handbuch der Physik, Vol. III/3. Springer-Verlag, Berlin (1965).
4. M. Slemrod, Admissibility criteria for propagating phase boundaries in a van der Waals fluid. Archive Rational Mechanics and Analysis, 81, 301-315 (1983).
5. R. Hagan, M. Slemrod, Viscosity-capillarity admissibility criterion with applications to shock and phase transitions. Archive Rational Mechanics and Analysis, 83, 333-361 (1983).
6. M. Slemrod, Dynamic phase transitions in a van der Waals fluid. Univ. of Wisconsin, Mathematics Research Center, Summary Report #2298, (1981).
7. Gurtin, M., Thermodynamics and the possibility of spacial interaction in elastic materials. Archive Rational Mechanics and Analysis 19, 339-352 (1965).
8. J. E. Dunn, J. Serrin, On the thermomechanics of interstitial working, Univ. of Minnesota, Institute for Mathematics and its Applications, IMA Preprint #24 (1983).
9. E. C. Aifantis, J. Serrin, The mechanical theory of fluid interfaces and Maxwell's rule. Technical Report, Corrosion Research Center, Univ. of Minnesota (1980).
10. R. Hagan, J. Serrin, One dimensional shock layers in van der Waals type fluids. In preparation.
11. R. L. Pego, Nonexistence of a shock layer in gas dynamics with a nonconvex equation of state. Univ. of Wisconsin, Mathematics Research Center, Summary Report #2505 (1983).
12. J. K. Hale, Ordinary differential equations, second edition. Krieger Publishing Co., Huntington, NY (1980).

13. G. Dettleff, P. A. Thomson, G. E. A. Meier, H.-D. Speckman, An experimental study of liquefaction shock waves. *Journal of Fluid Mechanics* 95, 279-304 (1979).
14. D. Gilbay, The existence and limit behavior of the one-dimensional shock layer. *American J. of Mathematics*, 7, 256-274 (1951).

RH/JS/ed

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER 2666	2. GOVT ACCESSION NO. AD-A241 708	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) DYNAMIC CHANGES OF PHASE IN A VAN DER WAALS FLUID		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) R. Hagan and J. Serrin		8. CONTRACT OR GRANT NUMBER(s) DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of Wisconsin 610 Walnut Street Madison, Wisconsin 53706		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 2 - Physical Mathematics
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P. O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE March 1984
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		13. NUMBER OF PAGES 28
		15. SECURITY CLASS. (of this report) UNCLASSIFIED
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) phase transitions, shock layer, van der Waals fluid, interstitial working, Liapunov function		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) This paper gives sufficient conditions to guarantee the existence of a shock layer solution connecting two different equilibrium states in a van der Waals fluid. In particular, the equilibrium states can belong to two different phases of the fluid. The constitutive laws come from a modified Korteweg theory which is compatible with the Clausius Duhem inequality. The Clausius Duhem inequality in turn gives rise to a Liapunov function. The main mathematical tool is the LaSalle invariance principle.		